

THEORETICAL MODELLING OF NANODEVICES USING EMBEDDED MOLECULAR CLUSTER METHOD

E K SHIDLOVSKAYA

*Information Systems Management Institute
Department of Natural Sciences and Computer Technologies
Lomonosova iela 1, k-6, LV-1019, Riga, Latvia
e-mail: shidlovskaya@inbox.lv*

ABSTRACT

Applicability of cluster embedding method with non-orthogonal wave functions for theoretical study of processes in nanodevices is studied. Processes in nanodevices are treated in the frameworks of time-dependent DFT. We demonstrate that our cluster embedding method is compatible with DFT Kohn-Sham method and quantum transport theory based on time-dependent DFT. We conclude that quantum transport theory approach for electric current calculation may be applied for non-orthogonal wave functions. Possibilities of theoretical modelling of nanodevices from the first principles are discussed.

Keywords: embedded molecular cluster model, non-orthogonal wave functions, time-dependent DFT, quantum transport theory, current in nanodevices

1 GENERAL

When we theoretically describe nanodevice we have to treat the whole quantum system as two subsystems: small finite fragment of the system containing nanodevice (cluster) and the rest of the system containing electrodes. Problem "cluster in the field of the rest of system" is successfully solved in the frameworks of embedded molecular cluster (EMC) model with *orthogonal* wave functions. We have modified EMC model treating cluster embedding problem in the frameworks of one-electron approximation with *non-orthogonal* wave functions. We have proposed new cluster embedding scheme based on our approach [1].

Our present aim is application of our cluster embedding method for quantum-chemical modelling of processes in nanosystems and calculation of electrical properties of nanodevices. One of the approaches for theoretical description of nanodevices is quantum transport theory developed by Gross with co-workers [2]. We study possibility to combine our approach with approach of Gross et al [2] based on time-dependent DFT (TDDFT). We demonstrate [3] that our cluster embedding method is compatible with DFT Kohn-Sham method. We conclude that our embedding scheme may be combined with TDDFT if electron transitions are described correctly: occupied and vacant cluster states are localized in the cluster region in the same manner. To get occupied and vacant states of the same localization degree, we have modified [3] our initial cluster embedding equations [1]. We demonstrate that our cluster embedding method is compatible with electric current calculation method based on TDDFT [2] and propose approach for calculation of electric parameters of nanodevices. Possibilities of our approach for theoretical modelling of nanodevices from the first principles are discussed.

REFERENCES

- [1] Shidlovskaya E K 2002 Improved embedded molecular cluster model. *Int. J. Quantum Chem.* **89** 349-370
- [2] Kurth S, Stefanucci G, Almladh C-O, Rubio A, Gross E K U 2005 Time-dependent quantum transport: A practical scheme using density functional theory *Phys. Rev. B* doi: 10.1103/PhysRevB.72.035308
- [3] Shidlovskaya E K 2012 Cluster embedding method with non-orthogonal wave functions for simulation of nanodevices *Nanodevices and Nanomaterials for Ecological Security* NATO Science for Peace and Security Series B: Physics and Biophysics, eds. Y.N. Shunin and A.E. Kiv, Springer 2012 pp. 191-202