

**LATVIJAS UNIVERSITĀTES
75. STARPTAUTISKĀ KONFERENCE**

ĶĪMIJAS SEKCIJA

Tēžu krājums

2017


N-SULFONYLCARBOXAMIDE AS AN OXIDIZING DIRECTING GROUP FOR RUTHENIUM CATALYZED C-H ACTIVATION/ ANNULATION

Elina Petrova

Latvian Institute of Organic Synthesis, Aizkraukles 21, Riga, LV-1006, Latvia
e-mail: petrova_eliina@inbox.lv

Functionalization via C-H activation enables atom and step economic synthesis of the complex molecules and has motivated a considerable progress of method development in recent years [1]. An effective C-H activation methodology has provided new routes for the construction of the variety of heterocycles. Isoquinolone derivatives can be obtained by Ru catalyzed C(sp²)-H activation of amides followed by the annulation of the intermediate metallacycle with olefins or alkynes [2].

We have developed an efficient traceless method for the synthesis of isoquinolones by Ru catalyzed annulation of *N*-sulfonylcarboxamide with alkyne. In this reaction, *N*-sulfonylcarboxamide acts as both an internal oxidant and directing group. Catalyst re-oxidation takes place in the N-S bond cleavage step. Of all *N*-sulfonylcarboxamides studied, the most efficient was *N*-2,6-difluorobenzene-sulfonamide which led to the formation of unstable sulfinate and readily decomposed to 1,3-difluorobenzene under the reaction conditions.



Supervisors: Dr. chem. Aigars Jirgensons, Dr. chem. Dace Rasiņa

References:

- [1] For the selected recent reviews see: (a) Gensch, T.; Hopkinson, M. N.; Glorius, F.; Wencel-Delord, J. *Chem. Soc. Rev.* **2016**, *45*, 2900; (b) Zhu, R.-Y.; Farmer, M. E.; Chen, Y.-Q.; Yu, J.-Q. *Angew. Chem. Int. Ed.* **2016**, *55*, 10578; (c) Davies, H. M. L.; Morton, D. *J. Org. Chem.* **2016**, *81*, 343. (d) Arockiam, P. B.; Bruneau, C.; Dixneuf, P. H. *Chem. Rev.* **2012**, *112*, 5879; (f) McMurray, L.; O'Hara, F.; Gaunt, M. J. *Chem. Soc. Rev.* **2011**, *40*, 1885; (g) Gutekunst, W. R.; Baran, P. S. *Chem. Soc. Rev.* **2011**, *40*, 1976. (h) Yamaguchi, J.; Yamaguchi, A. D.; Itami, K. *Angew. Chem. Int. Ed.* **2012**, *51*, 896; (i) Ackermann, L. *Chem. Rev.* **2011**, *111*, 1315.
- [2] (a) Ackermann, L.; Lygin, A. V.; Hofmann, N. *Angew. Chem. Int. Ed.* **2011**, *50*, 6379; (b) Ackermann, L.; Fenner, S. *Org. Lett.* **2011**, *13*, 6548; (c) Li, B.; Feng, H.; Xu, S.; Wang, B. *Chem. – Eur. J.* **2011**, *17*, 12573; (d) Li, B.; Ma, J.; Wang, N.; Feng, H.; Xu, S.; Wang, B. *Org. Lett.* **2012**, *14*, 736; (e) Yang, F.; Ackermann, L. *J. Org. Chem.* **2014**, *79*, 12070; (f) Krieger, J.-P.; Ricci, G.; Lesuisse, D.; Meyer, C.; Cossy, J. *Chem. – Eur. J.* **2016**, *22*, 13469; (g) Allu, S.; Swamy, K. C. K. *J. Org. Chem.* **2014**, *79*, 3963.